

Focus Review Article

Quorum Sensing Inhibitors: Current Progress of Natural Antimicrobials

Mohammad Asif

Department of Pharmacy, GRD (PG) IMT, Dehradun, (Uttarakhand), 248009, India


Receive Date: 11 March 2019, Revise Date: 17 September 2019, Accept Date: 10 October 2020

Abstract:

Misuses of antimicrobials in infectious diseases have led to the progress of extensive resistance in the infectious organisms. The unsuccessful of accessible antimicrobials to control infections makes it essential to discover alternatives to currently existing drugs. Their connection to infectious diseases and their natural ability increase antimicrobial resistance in microbes, has led to a platforms for research focused on new techniques to control them. In this affords we studied the pathogenicity in many bacteria was regulated by Quorum sensing (QS). The inhibition of QS system may cause the reduction of virulence and protect against bacterial infections. These bacteria rely on chemical communication (or QS) to coordinate activities necessary for their survival in groups by some course of action. Their dependence on QS has made those signaling systems within bacteria an attractive target for the design of new anti-infective agents. Compounds that can interrupt these processes are known as QS inhibitors. The QS is the key regulator of virulence in various bacteria. Various plants extracts and their chemical constituents were showed their effects on bacterial virulence factors by inhibiting of QS genes and QS-controlled factors and effects on bacterial growth. The anti-QS approach has promise in the fight against infections pathogens, thereby making the bacteria more susceptible to traditional antimicrobials. The QSIs may provide the newest weapon against infections involving drug-resistant bacterial strains. These QSIs come from a variety of sources and have a wide array of structures.

Key words: Quorum sensing inhibition; bacterial communication; antibacterial agents

Graphical Abstract:



Mohammad Asif was born in India. He studied Bachelor of Pharmacy (Pharmaceutical Chemistry) from IFTM, Moradabad, affiliated with Rohilkhand University Bareilly (U.P) in year 2003. Received the M.S. degree in Pharmaceutical Chemistry at Bundelkhand University Jhansi (U.P) in 2006 and the Ph.D degree also in Pharmaceutical Chemistry at Uttarakhand Technical University, Dehradun in year 2015. He focused his doctoral thesis on the Synthesis and biological evaluation of some new pyridazine-3(2H)-one derivatives. His research interests focus on the Medicinal Chemistry, Inorganic Chemistry, Organic Chemistry, Chemistry of Natural Products, Pharmaceutical Analysis and Physical Chemistry.

Introduction

*Corresponding author: Mohammad Asif Email: aasif321@gmail.com



Recently, microbial quorum sensing (QS) has come out as a new technology, because microbial population occupy a limited space, concentrations of extracellular signalling molecules accumulate, providing stimulus for unique cellular responses and protections from competing microbial communities [1, 2]. The eukaryotic organism produces metabolites that interfere with bacterial communication. *Delisea pulchra* produces a range of metabolites known as halogenated furanones. These compounds exhibit a broad range of biological activities, including antimicrobial activities [3]. Some polyphenolics having a gallic acid moiety (epigallocatechin gallate, ellagic acid and tannic acid) are commonly present in various plant species that inhibited QS communication between bacteria [4-6]. These polyphenolic depositions were correlated with a reduction of bacterial multiplications. The QS and the wide range of biological functions of QS systems is a targets for new antimicrobial agents [7-10]. QS-inhibitors, acting alone or with traditional drugs, could provide the newer generation of antimicrobial agents [11, 12]. The stimulating of QS systems has been responsible for a variety of physiological behavior in the bacteria such as bioluminescence, release of virulence factors, production of antimicrobials and biofilm formation [13-15]. This performance is generally observed to be controlled by cell density, but other conditions such as nutritional and environmental conditions can affect QS activities [16].

The QS system occurs in many bacterial species across both gram-positive and gram-negative bacteria. Bacteria liberate regulatory chemicals, called autoinducers (AIs), into their local environment [17]. As the population increases, concentration of the AIs also increases. At given concentration thresholds, the AIs bind to receptors in nearby bacteria, thereby setting in motion the QS cascade. Three major types of AIs have been identified: acylated homoserine lactones (AHLs) in gram-negative bacteria, auto-inducing peptides (AIPs) in gram-positive and autoinducer-2 (AI-2) molecules in both gram-positive and gram-negative bacteria. The different classes of AIs often target different receptors within the QS systems. AHLs target the LuxR family of transcriptional regulators, with the resulting AHL-LuxR-complex binding a variety of promoters and activating transcription factors [18, 19]. The receptors for AIPs are integral membrane proteins in the bacterial membrane. These proteins contain a protein kinase domain, which upon activation by AIP binding undergoes auto-phosphorylation. The activated receptor phosphorylates a response regulator, which activates the specific genes. The AI-2 signal molecules are more

species dependent and target a variety of Lux receptors [20, 21].

Target areas for anti-quorum sensing activity

The drug to target the QS system of bacteria is generally divided in three main target areas:

- destruction of the signal molecule,
- inhibition of signal production, and
- Inhibition of the signal receptor.

However, the last two target approaches involve, drug molecules interrupt some step in the natural QS-signaling pathway. Recently, few examples of QSIs which inhibit QS-signal molecule production have been reported in various bacterial strains. These QSIs may be similar in structure to the natural ligand or different, either binding directly to the signal-binding pocket or elsewhere on the receptors, in either case preventing the resulting QS-signal [22, 23]. The QSIs can come in a wide variety of chemical structures and be obtained from different sources. The recent advances of anti-QS activity is new and promising antimicrobial agents [24, 25].

Unexploited potential for antimicrobial activity

Plants should be a great source of antimicrobial agents. In fact, about 75% of the antibiotics are derived from the natural source like actinomycetes, a specific group of Gram-positive bacteria [26]. Antibiotic-producing microbes possess genes which protect them from the toxic effects of these compounds. It is very easy for one microbe to acquire antibiotic resistance from another through plasmid transfer or transposons. Plants are genetically dissimilar from the organisms they are trying to eradicate. Thus, there is little chance for a microbe to gain resistance from a plant. Plants can produce a multitude of diverse antimicrobial compounds such as simple phenolics, polyphenolics, catechins, quinones, flavanones, alkaloids, and terpenoids. Like microbial antibiotics, these compounds are targeted at killing the pathogen and work via a non-species specific mechanism such as disrupting microbial cell membranes. However, plants have another way of dealing with microbes-targeting a cell's communication system [27]. One form of intercellular communication in bacteria is known as QS [28]. Breakdown of QS system causes an attenuation of microbial pathogenicity [29, 30]. The discovery of anti-QS agents in plants provides us with yet another type of antimicrobial agents.

Functions of quorum sensing

A classic approach to solving problems is to break a system under consideration down into functional

elements. For many organisms of interest within biotechnology, the role of QS is complicated and unobvious. The QS is believed to regulate competence development, sporulation, antibiotic synthesis, virulence factor induction, cell differentiation, and nutrient flux along with other physiological events in pathogenic bacterial infections [31-33]. The QS was linked through proteomic analysis to increased pathogenic competence in tubercular strains of *P. aeruginosa* [34]. As biofilms age, cellular differentiation and death enhance nutrient sequestration and allow for biofilm sustenance when nutrients become scarce [35,36]. Many human or plant pathogens exist as biofilms in their hosts, but occupy a different modality both pre- and postinfection. QS is thought to regulate this transition. For example, in the human pathogen *V. cholera* quorum signalling enables the cells to negotiate the acidic human gut environment without compromising their ability to infect. Biofilms formed by *V. cholerae* protect the cells from stomach bile and low pH [36,37]. In addition to regulating intraspecies survival and differentiation in bacterial communities, QS also relates interspecies information between symbionts and competitors [38, 39]. Acyl-homoserine lactone (AHL) signals from *P. aeruginosa* can enter mammalian cells and activate artificial transcription factors, although it is yet unclear which native eukaryotic proteins may be transcribed through this mechanism [40, 41].

Quorum sensing a novel target

Quorum sensing (QS) is a population-dependent phenomenon and its ability to sense the size of a bacterial population mediated through small signaling molecules or autoinducers (AIs) [28]. These AI molecules are constantly produced and received at a basal level by bacterial cells. With high population density, there is an excess of signaling AI molecules in their environment. These signals diffuse back into the cell where they facilitate the regulation of gene expression. QS-systems are everywhere among bacteria, and have found to regulate diverse functions such as luminescence, biofilm formation, antibiotic and virulence factor generation, pigment production, plant-microbe interactions, and motility [43]. Although there are different QS systems [44], the most widely considered model is based on the Lux system of *Vibrio fischeri* and *V. harveyi*. This QS phenomenon involves a three component-system: a freely diffusible signal, a synthetase to make this signal, and a regulator that interacts in conjunction with the signal to regulate gene expression. The main signaling molecules produced by Gram-negative bacteria are AHLs. They differ in the length of their side chains and substitution, based on the organism that produces those [45]. In *V. fischeri*,

LuxI produces an AHL signaling molecule which binds to LuxR at a definite concentration. This concentration is reached only when adequate bacteria are present to produce satisfactory amounts of AHL. The *V. fischeri* LuxR-AHL-complex can then bind to the operator and up regulate transcription of the luciferase genes [46-49]. The AHLs bind to LuxR (product of *luxR*) at a certain concentration and activate transcription of *luxI* and the luciferase genes [44]. AHL-mediated QS-systems based on the LuxI/R model have been described in human pathogens like *P. aeruginosa*, *Yersinia pseudotuberculosis*, and *E. coli*, as well as plant associated bacteria like *Rhizobium leguminosarum*, *Ralstonia solanacearum*, and *Erwinia carotovora* [50]. In all cases QS- systems may regulate virulence. The QS has given a new target to attack and attenuate bacterial pathogenicity [51-55].

Mechanisms of QS inhibition

There are a number of ways to inhibit cell-cell communication (or QS) including competitive inhibition, signal binding, degradation of the signaling molecule, and inhibition of genetic regulation systems. The success with competitive inhibition, furanones and many other QSIs has been discovered [56]. These QSIs are based on the C12-AHL structure and cause a reduction in LasR activity. AHL-antibodies have been developed to suppress QS through signal binding [57, 58]. A C12-AHL-protein conjugate was able to successfully inhibit *lasB* expression genes, and a similar molecule with extremely high binding affinity for C12-AHL. Blocking S-adenosyl methionine (SAM) or the fatty acid precursors necessary to produce AHLs leads to decreased production of C12-AHL by LasI [58, 59]. Many bacteria including *Bacillus* sp, *Variovorax paradoxus*, *Arthrobacter* sp, and *A. tumefaciens* produce lactonases, enzymes that cleave and deactivate the lactone ring of various AHLs [60, 61]. Lactonase expression in *P. aeruginosa* significantly decreases AHL production and virulence factor expression. The practical applications of anti-QS especially in drug resistant bacteria therapy because of increased incidence of drug failure due to the large number of pathogenic bacteria developing resistance to available antimicrobials [62].

Linking traditional and modern medicine

The enormous numbers of plants with chemical diversity are the basis for medicinal compounds [63]. However, this inherent quality of plants helps to search for new drug molecules. Most plants contain constitutive or inducible defense against phytopathogenic bacteria. Plants used medicinally because they contain chemical defenses specific to human pathogens [64,

65]. However, ethnopharmacology is beneficial to traditional societies. The chemical and pharmacological activities of the plants are potentially useful and unique medicinal values. Consequently, most studies focus on ethnopharmacology, depending on the expertise and interests of the researchers [66].

Ethnobotany for looking to the future antimicrobial agents

Although synthetic drugs can be provide pharmacological effects to cure diseases. The high biological diversity is linked to a high chemical diversity that allowing for the evolution of numerous bioactive natural molecules. Many plant and herbal drug molecules already exist in nature and many are waiting for discovery [67-69]. Plants and herbs create specific compounds to serve in reproduction and defense mechanisms [70]. Plants and herbs are having toxic and bioactive compounds to protect them against herbivore and pathogen attacks. Because plants and herbs are rely on chemical defenses. Various plants or herbs and their fruits were shown anti-QS activity in a *C. violaceum* biomonitor strain and on the swarming motility of *E. coli* and *P. aeruginosa* [71]. Fruits including raspberry, blueberry, blackberry, cranberry, and grape, and herbs such as ginger, kale, oregano and turmeric exhibited moderate inhibition of QS processes. However, no compound was obtained in animal as QSIs. The furanones and their synthetic derivatives, the compounds ellagic acid, tannic acid, and epigallocatechin gallate have been shown to inhibit QS in both an *E. coli* and *P. aeruginosa* biomonitor strain [72]. The related polyphenolics should be further explored as anti-QS compounds.

Anti-quorum sensing and antibacterial from ethnobotanicals

Many research based on ethno-botanical for antibacterial activity for demonstrating not only the need for antimicrobial drugs, but also the large number of plant species utilized for bacterial pathogenicities [73]. Although many medicinal plants and herbs base research has led to the discovery of many important medicines, there is not a more overlap between commonly used plant medicines and antimicrobials [74-78]. However, there are some plants that have made their way to the drug developments. The discovery of some natural compounds that inhibit cell-to-cell communication (QS), could provide a novel method of combating pathogenic infections [79]. Anti-QS agents were first characterized from red marine alga, *Delisea pulchura*. This alga was showed anti-fouling properties, and was found to contain halogenated furanones which block AHLs via

competitive inhibition and destabilization of LuxR [80]. The structural similarity of furanones allows to competitively inhibit the action of AHL signaling molecules (**1a** and **1b**) (fig 1) [81]




Figure 1. Acyl-homoserine lactones signaling molecules

Various naturally occurring phyto-molecules, particularly extracts from plants and dietary substances, have been evaluated to modulate LuxR-type QS in Gram-negative bacteria. However, mechanism of action of many of these compounds is poorly understood. A significant deviation from the AHL framework is usually observed and there is little structural correlation with any other anti-QS agents whose molecular target is known. For example, the some natural substances are known to modulate various AHL-mediated QS systems or AHL-regulated phenotypes are clove oil (*C. Violaceum* and *P. aeruginosa*), dietary phytochemicals (secondary metabolites of plants, *C. Violaceum* and *P. aeruginosa*) [71,82], honeys (*E. carotovora*, *Yersinia enterocolitica*, *Aeromonas hydrophilia*, and *C. Violaceum*), cranberry juice (*V. harveyi*), extracts of *Tremella fuciformis* (white jelly mushroom, *C. Violaceum*), extracts of some medicinal plants from the Indian subcontinent (*P. aeruginosa*), extracts of pea (*Pisum sativum*) seedlings (*C. Violaceum*) [83], extracts of *Medicago truncatula* seedlings (*P. putida* CepR reporter, *E. coli* LuxR reporter, *E. coli* LasR reporter, and *C. Violaceum*) [84] and extracts of South Florida plants (*P. aeruginosa*) [85].

Anti-QS activity of several plant extracts have been reported recently, and revealed that the QSI organism protect themselves from invasion of other pathogenic microbes. Plant, bacteria, fungi could produce compound which could interfere the QS-regulated gene expression in the invading organism [86,87]. Since the discovery of AHL inhibitors in *D. pulchura*, anti-QS activity and South Florida *Caulerpa* species and various higher plants including various fruits and vegetables [83,88]. *Pisum sativum* (pea) seedlings and root extracts produced an inhibition of pigment production, exochitinase activity, and protease activity in *C. violaceum* [71]. Garlic (*Allium sativum*), carrot (*Daucus carota*), chamomile (*Matricaria* sp.), water lily (*Nymphaea* sp.) and some peppers (*Capsicum* spp.) were found to possess anti-QS activity against a *luxI-gfp* reporter strain. Toluene extract of *Allium sativum* or garlic exhibited anti-QS activity against Gram-negative transcriptional regulators Lux R or Las R [87]. Garlic was also inhibited biofilm formation and prevented nematode death [23]. Not only garlic but also several other plants extract like *Thymus vulgaris* family

alone had little to no effect on the biofilm [87], compounds **3** and **4** all were identified as LuxR antagonists, most significantly cyclic thioacetal **4a** and cyclic disulfide **4b** showed only anti-QS activity, while **3a-d** were also toxic to the bacteria. The *in-vivo* used of garlic extract as a potential therapy for lung infections [108]. Mice infected with *P. aeruginosa* were treated with garlic and tobramycin combination showed significant clearing of the bacterial infections as compared to control group. Through bioassay-guided fractionation, six sulfur-containing compounds from the garlic extract, **3a-d** and **4a-b**, were identified that showed anti-QS in a LuxR monitor system [87]. Compounds **3a-d** antagonized LuxR but were toxic to the bacteria. Compounds **4a** and **4b** possessed anti-QS activity exclusively, but only in the LuxR monitor system; none of the sulfur containing compounds **4a** and **4b** have active against *P. aeruginosa* QS. A number of AHL derivatives that incorporated sulfide, sulfinyl, and sulfonyl motifs were found to inhibit either one or both of the LuxR and LasR-QS systems.

Alkaloids from natural sources, recently reported as a natural anti-QS alkaloidal agents, solenopsin A (**5**, Fig. **9**) [109]. A toxic alkaloid from the fire ant *Solenopsis invicta*, solenopsin A has showed an inhibitor of

phosphatidylinositol-3-kinase signaling and angiogenesis [110] as well as a potent cardiorespiratory depressant [111]. In *P. aeruginosa*, solenopsin A inhibits pyocyanin production, an indication of QS signaling suppression. Synthetic AHLs indicated that the rhl signaling system might be the target of solenopsin. Solenopsin A (**5**), a venom alkaloid from the fire ant *Solenopsis invicta*, inhibits QS in *P. aeruginosa* [109]. This compound is structurally reminiscent of OdDHL (natural AHL of the LasR system) in that both contain a long hydrocarbon chain attached to a N-containing heterocycle, via a chiral-C; however, both structures contain distinct molecular frameworks. Interestingly, exogenously added BHL, but not OdDHL, restored *P. aeruginosa* QS signaling, suggesting that solenopsin A actually targets the BHL-dependent *rhl* QS system. Solenopsin analogues **5a-e**, (Fig. **4**), no analogue demonstrated to increased anti-QS activity relative to the parent compound. Analogues of **5** with shorter acyl chains, however none showed any increased QSI activity. Further investigations showed that solenopsin A reduced biofilm production in *P. aeruginosa* in a dose dependent manner, indicating a QS signaling suppression mechanism.




Figure 3. QSIs from garlic extract.




Figure 4. Alkaloid QSI, solenopsin A, isolated from *Solenopsis invicta*.

Essential oils from various plants can affect biofilm formation and structure, among these cinnamaldehyde (**6a**) and its derivatives (Fig. **5**). The cinnamaldehyde showed *E. coli* biofilms inhibitor. The connection of cinnamaldehyde to QS was first demonstrated by Gilbert and co-workers [112,113], who showed the compounds ability to inhibit both AI-1 and AI-2 receptor systems. The bioluminescence of the 3-oxo-C4-HSL mediated *V.harveyi* BB886, as well as the AI-2 responsive reporter strain *V.harveyi* BB170 were significantly inhibited at 60 and 100 $\mu\text{mol mL}^{-1}$ respectively. The effects of cinnamaldehyde and substituted cinnamaldehyde derivatives on AI-2 reporter systems by examine the bioluminescence of several *V. harveyi* mutants. The two most potent compounds were cinnamaldehyde and 2-nitrocinnamaldehyde (**6b**). The inhibitors affected the

target protein LuxR and further studies with these compounds showed biofilms inhibitory effect on two vibrio mutants, *V. anguillarum* 4411 and *V. vulnificus* LMG [114]. The **6a** affected the total mass of the biofilm, but not the number of viable cells. These compounds were affecting the production/accumulation of the exopolysaccharide matrix. Plant essential oil components and isolated those that modulated biofilm formation in *E. coli* [112,113].

One of these components was identified as cinnamaldehyde (**6a**), which was reported as an inhibitor of 3-oxo-C6-HSL (OHHL) QS in *E. coli* and 3-hydroxy-C4-HSL QS in *V. harveyi* at sub inhibitory concentrations (Figure 5) [112]. In the case of *V. harveyi*, inhibition was not selective for AHL QS, with the AI-2 system also being affected. Interestingly,

cinnamaldehyde (**6a**) had less effect on *lasR* promoter activity, induced by OdDHL, in an *E. coli* strain containing a LasR biosensor (OdDHL inducible). The 3-C aliphatic side chain of cinnamaldehyde (**6a**) may interfere with the binding of the smaller 3-OH-C4- and 3-oxo-C6-HSL AHLs to their cognate receptors but was not sufficiently long enough to reduce the binding of OdDHL to LasR. Recently, *p*-coumaric acid (**6c**), has putative antagonistic activity against *Chromobacterium*, *Agrobacterium*, and *Pseudomonas* QS. *p*-Coumaric acid (**6c**) and cinnamaldehyde (**6a**) share a common structural motif, namely, an R-unsaturated carbonyl functionality connected to a phenyl ring system. Recently, extracts of the bark of *Combretum albiflorum* for their capacity to inhibit the production of extracellular virulence factors in *P. aeruginosa*.




Figure 5. Cinnamaldehyde (**6a**), 2-nitrocinnamaldehyde (**6b**) and *p*-Coumaric acid (**6c**).

Extract from other plant species have been shown to inhibit bacterial QS. While the specific anti-QS compounds from these extracts were not reported, so further investigation is necessary to identify their active compounds. The QSI activity by vanilla extract in the mutant *Chromobacterium violaceum* CV026 which lack the autoinducer synthase [115]. The inhibition of QS-controlled virulence factors in *P. aeruginosa* by extracts from several South Florida plants [116]. Three of these extracts, those from *Conocarpus erectus*, *Bucida buceras* and *Callistemon viminalis*, showed inhibition of *LasA* protease, *LasB* elastase, pyoverdinin production and biofilm production.

Polyphenolic QSI Compounds

Polyphenols are well known antioxidant, and have been shown to effect on biofilms. The connection between polyphenols and QS is established [117] by both biofilm and QS inhibition for such compounds (-)-epigallocatechin (**7**, EGCG), ellagic acid (**8**) and tannic acid (**9**) (Fig. (6)). by using two AHL-dependent QS strains *P. putida* (pKR-C12) and *E. coli* MT102, and identified as antagonistic effect. EGCG was the most active followed by ellagic acid, then tannic acid. Biofilm inhibition was showed against in *B. cepacia*. Tannic acid showed little reduction in film thickness, both EGCG and ellagic produced a marked reduction, with ellagic acid being the most effective. Several other polyphenols have shown promise as anti-QS activity (Fig. 7). The antagonistic effect of pyrogallol (**10**) analogues was on QS systems in *V. harveyi* [118], a different approach to screening compounds for potential anti-QS activity was reported [119]. Using the crystal structure of TraR from *A. tumefaciens*, various compounds from TCM with antibacterial activities were compared to known QSIs, furanone C30. Some compounds showed docking scores higher than that of furanone C30. Of those, few compounds were shown to inhibit biofilm formation in *P. aeruginosa*, the most potent being the flavanoid baicalein (**11**). In addition, **11** showed to have a marked synergistic effect, reducing biofilm formation when used in combination with the antibiotic, ampicillin. Baicalein was also shown to promote proteolysis of the receptor TraR protein in *E. coli*. Rosmarinic acid (**12**), from the root of sweet basil, is a caffeic acid ester showed inhibitory growth of some *Pseudomonas* species and interfere with QS activity [120]. Several active fractions containing flavonoid-like compounds; one active compounds flavan-3-ol catechin (**13**), which was also found to have a negative impact on the transcription of several other QS-related genes in this bacterium. Catechin (**13**) is thought to possibly interfere with the perception of the native AHL by the LuxR homologue RhlR, though little is known about the precise mechanism of action [87].




Figure 6. Polyphenolic QSIs, (-)-epigallocatechin (**7**), ellagic acid (**8**) and tannic acid (**9**).




Figure 7. Polyphenolic QSIs, pyrogallol (10) and baicalein (11) and Rosmarinic acid (12) Catechin (13).




Figure 8. Polyphenolic QSIs, pyrogallol (10) and baicalein (11) and Rosmarinic acid (12) Catechin (13).

Polypeptide QSI Compounds

The signaling molecules in QS systems of gram-positive bacteria are often polypeptides. One of the most extensively investigated anti-QS peptides has been RNAIII inhibiting peptide (RIP; YSPWTNF-NH₂). RIP interferes with the QSQ1 system in gram-positive bacteria. QSQ1 consists of autoinducer-RNAIII-activating protein (RAP) and its target molecule TRAP. RIP competes with RAP for TRAP binding. Consequently TRAP phosphorylation is inhibited and agr expression is reduced. This QS system has potential target for new antibacterial agents, particularly against resistant strains [121]. Evidence has shown RIP to be effective against *S. aureus* by the disruption of QS mechanisms [122,123]. Numerous researches have showed specific *in-vivo* evidence for biofilm inhibition in *S. aureus* and *S. epidermidis* [124-128]. The use of polymethylmethacrylate beads loaded with RIP has also showed an effective system for treatment of drug resistant film from orthopedic pathogens [129]. RIP has also showed synergistic with a variety of antibiotics drug in eliminating biofilms from both *S. aureus* and *S. epidermidis* [130-133]. Considerably, SQS2 sensing inhibitors, which rely on agr-encoded autotinducing peptides (AIPs), while several groups have reported the inhibitory AIPs [134-138].

QSI Antibiotics

Antibiotics are the first choice for many microbial infections, in many cases the lack of bactericidal activity caused them useless. In chronic *P. aeruginosa* infections of cystic fibrosis, macrolides such as azithromycin (14, AZM) are not often included in the treatment regimen. However, more evidence that low (sub-MIC) doses of a number of antibiotics (Fig. (8)

which may elicit a variety of other effects. The AZM is to interfere with QS regulation in *P. aeruginosa* PAO1 [139]. The azithromycin interferes with transcription of AI synthase. Investigations reported that some different antibiotics for QSI activity [140]. The screens on QSIS1 revealed three compounds with strong QSI activity, such as ceftazidime (15, CFT) and ciprofloxacin (16, CPR), as well as the AZM. The direct connection between QSI effects and biofilm formation showed that AZM reduced the adherence of *P. aeruginosa* to polystyrene at low concentrations (1-2 mg/L). The connection between biofilm inhibition and inhibition of AI synthesis [141] concluded that the biofilm inhibition was at least partially due to interference of both C4-HSL and 3-oxo-C12-HSL formation. This would reduce the production of extracellular virulence factors which is necessary for biofilm formation. The *in-vivo* effect of AZM [142] of chronic *P. aeruginosa* lung infections gave significant improvement of clearance of alginate biofilms and reduced the severity of lung pathology. Presumably, due a suppression of QS-regulated virulence facts, in particular LasB expression, thus impairing the bacteria's ability to produce fully formed the biofilms.

QSI Compounds from fungi

Anti-QS agents were first described in the red marine alga *Delisea pulchra* [3], in a south Florida alga and a few higher plants [84,108]. In search for QSIs in some medicinal plants, it has been shown that terrestrial plants not only produce autoinducer [143] increases to confound the bacterial QS system [116,144]. The anti-QS medicinal plants, efficient screenings confirmed some plants, *Conocarpus erectus*, *Chamaesyce hypericifolia*, *Callistemon viminalis*, *Bucida buceras*, *Tetrazygia bicolor*, and *Quercus virginiana*, have anti-QS properties using *Chromobacterium violaceum* and

Agrobacterium tumefaciens NTL4 strains as biomonitors. These plants showed potentially therapeutic effects against bacterial infections (Bauer and Mathesius 2004). The fungal metabolites that interfere with bacterial QS signaling [23] reported the screening of various penicillium species. The initial screen used the QSI1 selector, which identifies lux inhibitors, with a secondary screen using QSI2 selector for *P. aeruginosa* QS systems. Various extracts were showed QSI effects. Two of these secondary metabolites were penicillic acid (**17**) and patulin (**18**) (Fig 9). Both were showed QS-controlled gene expression in *P. aeruginosa*, most likely affecting the RhlR and LasR proteins. *In vitro* studies showed that *P. aeruginosa* PAO1 biofilms treated with patulin were more susceptible to the antibiotic tobramycin. The *in vivo* study showed the use of patulin as a used for pulmonary infections with *P. aeruginosa*. The QSI showed a significantly lower mortality rate and faster lung clearance. These two compounds were effect on biofilms of *Klebsiella sp.*, *B. subtilis* and *Bacillus cereus* [145]. The study was to measure any interaction with LuxR AI-2 sensing and LuxS based AI-2 production, two receptors that could not be evaluated in the *P. aeruginosa* PAO1. Interestingly, penicillic acid alone had no effect on biofilms, however in combination with ethylenediaminetetracetic acid (EDTA) it enhanced biofilm formation in both *Klebsiella sp.* and *B. cereus*. Patulin alone promoted biofilm formation in *B. cereus* with a significant enhancement in both bacillus species when combined with EDTA.




Figure 9. QSI fungal metabolites, penicillic acid (**17**) and patulin (**18**)

Brominated furanones

Furanones are compounds which have substantial interest due to their considerable QSI activity of brominated furanones or fimbrolides. Natural furanones were isolated from red algae *Delisea pulchra*, which produces more than 30 of furanones as secondary metabolites. The algae use as protection against colonization from both prokaryotes and eukaryotes. These compounds were interfered with QS-controlled swarming phenomenon in *Serratia liquefaciens* and *Proteus mirabilis* [43]. These furanones interfering with AHL-mediated QS is extensive [3,80,146] with a number of compounds showing promise as QSIs (Fig. (10)).




Figure 10. Natural and non natural brominated furanones

One furanone isolated from *D. pulchra* showed direct effect on biofilm formation is (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2-(5H)-furanone (**19**) [147]. Compound **19** inhibited biofilm formation of *E. coli* XL1-blue on mild steel surfaces with little toxicity to the bacteria. Further studies [148] showed, with a variety of *E. coli* mutants, point to the inhibition of the AI-2 system as the target for the furanone. Further evidence for AI-2 interaction was reported in *Bacillus subtilis*, where **19** reduced biofilms significantly at 40 $\mu\text{g mL}^{-1}$ [149] and using full-genome DNA microarrays showed **19** to induce 92 different genes in *B. subtilis* [150]. Similar growth inhibition was also shown in *Desulfotomaculum orientis* [30,150]. These compounds showed to disrupt AHL QS [62], these compounds increased the susceptibility of the bacteria to the antibiotics. In addition, biofilms grown in the presence of **20b** were readily dissolved overnight by sodium dodecyl sulfate. Furanone **21** had biofilm inhibition against the gram-positive bacteria, *S. epidermidis*, reducing biofilm formation by 68%, while having no effect on the growth of the bacteria [151].

Ellagitannin, a natural compound from various medicinal plants have shown anti-QS activity particularly against various Gram-negative bacteria [85,152]. Sub inhibitory concentrations of phenyl lactic acid, produced by *Lactobacillus* probiotic strains, have been shown to attenuate *P. aeruginosa* virulence and pathogenicity by interfering different regulated processes of QS. Tannic acid, a plant polyphenol, also showed anti-QS various Gram-negative bacteria. Ethanolic extracts of the plants *Sonchus oleraceus* and *Laurus nobilis* showed anti-QS in the Gram-negative bacterium *C. Violaceum* [90]. Vanillin extracts have anti-QS activity in *C. Violaceum*, [115] and vanillin (**22**) may be the active agent. Secondary metabolites of the North Sea bryozoan *Flustra foliacea* having a variety of brominated alkaloids, two of which (**23**, **24**) were found to specifically block AHL-regulated gene expression. Both compounds caused a reduction in the signal intensities in biosensor strains of *P. putidia* and *E. coli*. In addition, compound **23** was reduced the production of extracellular proteases in *P. aeruginosa*, a phenotype associated with the virulence of pathogens, which is under the stringent control of AHL-dependent QS systems. The extracts of the red alga *Ahnfeltiopsis flabelliformis* inhibited QS mediated by OHL and the TraR transcriptional activator protein [153]. Compounds betonicine (**25**), floridoside (**26**), and

isethionic acid (**27**), each of these compounds were showed QSI activities (Fig. 11). None of the three compounds exhibited inhibition when tested individually. In contrast, a complex of floridoside and isethionic acid revealed a dose-dependent inhibition on

OHL activity, these two compounds are responsible for the inhibition activity of red algae extract. Unexpectedly, betocine and *cis*-betocine showed a dose-dependent stimulatory effect in TraR-mediated QS responses.




Figure 11. Quorum sensing inhibitors (vanillin **22**), (alkaloids **23** and **24**) and (solenopsin A and analogues **25-27**).

A extracts from food sources and herbal medicines for QSIs and reported that 4-nitropyridine-*N*-oxide (**28**) and toluene extracts of garlic specifically inhibit QS-regulated gene expression in *P. aeruginosa*. Subsequent in vitro studies on *P. aeruginosa* biofilms demonstrated that garlic extract significantly reduced the tolerance of the bacteria to the antibiotic tobramycin. The garlic extracts can antagonize the activity of the QS receptors AhyR and TraR. Recently, garlic extract promotes rapid clearing of mice pulmonary *P. aeruginosa* infections [108].

Applications

Pathogen and pest (undesirable organism in environment) management comprise current applications of QS-technology. Inhibition of QS signaling is the most obvious, most omnipresent application of QS knowledge. The targeting of QS for the treatment of bacterial infections were reported in various studies [154,155]. Some examples using various aspects of QS are discussed. *Staphylococcal* subspecies, such as methicillin-resistant *S. aureus*, remain one of the most opportunistic pathogens. The QS inhibit the development of *S. aureus* and *S. epidermidis* biofilms. Both strains were susceptible to an RNA III inhibiting peptide (RIP), a specific QSI that interferes with the gene locus *agr*, which is responsible for *staphylococcal* toxicity [125]. When RIP was added to the culture, the cells were less able to form a biofilm. Expanding this approach to include antibiotics, the effect of combining RIP with seven different combinations of antibiotics and found that in every case the combination of RIP and an antibiotic greatly improved the performance of the antibiotic [125]. Looking to Gram-negative bacterial infections, *P. aeruginosa* from intubated patients revealed quorum-signalling targets that could be potentially inhibited to

prevent the formation of biofilms [156]. Most of the QS genes expressed were classified as transcription factors. High concentrations of AI-2 were detected in sputum from cystic fibrosis patients. The research found that *P. aeruginosa* transcriptional virulence factors could be stimulated by oral pharyngeal microflora and that high concentrations of AI-2 were present in in vitro cultures of both *P. aeruginosa* and the host microflora, indicating that the host's own bacteria might be stimulating the growth of *P. aeruginosa* [157]. Plants have long been known to interact with symbiont bacteria through QS signaling, and plant pathogens use QS signaling to colonise their hosts.

A quorum signaling in plant-pathogen symbiont interactions discusses some of the potential applications that could arise from these relationships [50]. *Erwinia carotovora* and *Agrobacterium tumefaciens* of AHL expression and repression in the plant rhizosphere [158]. Plant-microbe relationships with potential for pathogen control. *Burkholderia cepacia* in onions [159], *Ceratocystis ulmi* (a fungus that causes Dutch elm disease) [160], *Sinorhizobium melilot* (symbiont) and *P. aeruginosa* (pathogen) in legumes [161], and wine grape consortia [162]. Part of the biocontrol activity of *Bacillus thuringiensis* is through AHL lactonase, an AHL-degrading enzyme [163]. Biocontrol efficacy against *E. carotovora* was reduced in *B. thuringiensis* mutated for AHL lactonase. Also, strains of *E. coli* and *Bacillus fusiformis* lacking AHL lactonase showed a similar lack of anti-microbial capability when cultivated on potatoes [162]. Interestingly, *B. thuringiensis* did not inhibit the growth of *E. carotovora*, but rather, inhibited its virulence and ability to cause soft rot disease in potatoes. The virulence inhibition capability of *B. thuringiensis* might make it a candidate for fighting bacterial infections. The demonstrated the effect of

recombinant AHL lactonase in transforming strains incapable of biocontrol into biocontrol agents [164]. *P. fluorescens* was transformed with the *aiiA* gene encoding AHL lactonase under a constitutive promoter. AHL-degrading capability in-vitro and the ability to inhibit growth of aquatic biofilms [165] is another potential application of QSIs [166]. In a screen for fungicidal bacteria reported finding numerous strains capable of degrading invasive *Gymnodinium catenatum* [167]. Included in the fungicidal strains were *Pseudomonas* ssp. expressing AI-2; however, the occurrence of AI-2 did not correlate with fungicidal activity. In activated sewage sludge, QS has been shown to change the characteristics of waste water [168] and AHL compounds to activated sewage sludge and found an increase in the phenol degradative capacity. In a nonaquatic study of food spoilage, neither AI-2 nor AHL signalling correlated with biofilm formation of Gram negative bacteria [169]. The role of quorum signaling, especially with respect to its applicability to biological control, the problem with a wide variety of infections and their contribution to the antibiotic resistance of certain bacteria, have made controlling them an important facet in managing and removed bacterial infections. The structurally diverse groups of molecules have shown QSIs. By identifying active QSIs, various are natural QS compounds taken from different sources; the mechanism of QSI can be better understood. These QSIs hold great potential for the design of more effective antibacterial agents. The QS systems are regarded as new targets for antimicrobial strategies. In this perspective, it is important to consider the potential implication of horizontal transfers in the reacquisition of virulence.

Discussion

Applications for QS sensing are presently limited by perceptiveness of QS mechanisms. The use of QS, further perceptiveness of quorum functionality is required before the influence of this can be realized. The capability to block QS and makes the use of isolated compounds for driving protein expression. The full-scale treatment of the bacterial quorum circuit in a biotechnological purpose remains an unfulfilled goal. QS driven gene expression was used in concert with complicated regulation in eukaryotic cells to finely control product synthesis [170,171]. The method was effective in a rat model. The significance of the work is that it demonstrates a potential 'infection driven' gene therapy, where the existence of a bacterial extracellular signal may drive the expression of genes designed to fight the infection. Many bacteria utilize sophisticated regulatory systems to ensure that some functions are only expressed when a particular population density has reached [172-174]. The QS described form of density-dependent gene regulation which relies on the making and perception of small signal molecules by bacterial cells. As in many pathogenic bacteria the production of virulence factors is QS regulated, it has

been suggested that this form of gene regulation allows the bacteria to remain invisible to the defense systems of the host until the population is sufficiently large to successfully establish the infection. The polyphenolic compounds can interfere with bacterial QS. These polyphenols are widely distributed in the plants and may be important for promoting plant fitness.

Future perspectives

The use of small molecules to modulate bacterial QS systems has attracted significant interest over the course of the last 15 years. Various structurally diverse nonnative activators and inhibitors have been discovered; provide researchers with an expansive set of chemical tools to study this form of intercellular communication. Ultimately, strategies based upon the chemical modulation of bacterial QS may prove to be of value in a wide range of fields, including medicinal, agricultural, and environmental. However, chemotherapeutics purposes remain a long way off. Attaining a combination of efficacy *and* selectivity (small molecule modulation of a specific QS-regulated phenotype in a given bacterial species) presents a significant challenge [175,176]. There is a significant need for the standardization of the assays to study small-molecule modulation of QS-pathways. This facilitates the elucidation of more accurate SAR data for QS-modulators, which should enhance the perceptiveness of the molecular features crucial for desired biological activities. There is a definite need for more detailed fundamental studies into the molecular basis of QS modulation, that is, the mechanisms of action of small-molecule activators and inhibitors in terms of the fundamental bonding interactions involved. Such information would provide a scaffold for a deeper perceptiveness of the performance of existing small molecule modulators on a molecular level and also facilitate the rational de novo design of new next-generation agents with improved molecular properties (efficacy and selectivity). The field of small-molecule modulation of QS can be considered, in many regards, to still be in its infancy [177-182]. There is, thus, considerable scope for further exciting developments to be made in this area; the reliance of QS upon a language of small molecules undoubtedly means that chemists will play an integral role in such progress.

Conclusions

The plant kingdom has long been a source of diverse medicinal compounds, many ethnobotanically directed searches for agents that can be used to treat pathogenic infections, mainly focused on bactericidal effects. Our focus to anti-QS and anti-virulence properties may reveal quorum-quenching compounds from medicinal plants for provide a novel method for the treatment of infections against resistant strain of microorganism particularly against *P. aeruginosa*. The effects of the

medicinal plant extracts on *P. aeruginosa* are quite complicated and maybe extend beyond the domain of the QS control hypothesis. The failure of existing antibiotics to control infection makes it crucial to find alternatives to currently available drugs. The pathogenicity in many bacteria is regulated by QS that is the key regulator of virulence and biofilm formation in *P. aeruginosa* and other relevant bacteria. Various plants showed their effects on *P. aeruginosa* virulence factors and the QS system with significant inhibition of LasA protease, LasB elastase, pyoverdinin production, and biofilm formation. In fact, an anti-QS approach has already shown promise in the battle against *P. aeruginosa* infections. However, the reduction of QS gene expression and signaling molecule levels and the end effect on virulence factor production provide some insight into why these plants were used in the past and how they can be used in the future to combat *P. aeruginosa* and other bacterial infections.

References

- [1] J.T. Hodgkinson, M. Welch, D.R. Spring, Learning the Language of Bacteria. *ACS Chem. Biol*, 2(11) 2007, 715-17.
- [2] A. Jayaraman, T.K. Wood, Bacterial quorum sensing: signals, circuits, and implications for biofilms and disease. *Annu. Rev. Biomed. Eng*, 10 2008, 145-67.
- [3] M. Manefield, T.B. Rasmussen, M. Hentzer, J.B. Andersen, P. Steinberg, S. Kjelleberg, M. Givskov, Halogenated furanones inhibit quorum sensing through accelerated LuxR turnover. *Microbiology*, 148(4) 2002, 1119-27.
- [4] R.P. Ryan, J.M. Dow, Diffusible signals and interspecies communication in bacteria. *Microbiology*, 154(7) 2008, 1845-58.
- [5] C.D. Sifri, Quorum sensing: bacteria talk sense. *Clin. Infect. Dis*, 47(8) 2008, 1070-76.
- [6] P. Williams, K. Winzer, W.C. Chan, M. Camara, Look who's talking: communication and quorum sensing in the bacterial world. *Phil. Trans. R. Soc. Lond. B*, 362 2007, 1119-1134.
- [7] T. Bjarnsholt, M. Givskov, Quorum sensing inhibitory drugs as next generation antimicrobials: worth the effort? *Curr. Infect. Dis. Rep*, 10(1) 2008, 22-8.
- [8] B. Raffa Robert, R. Iannuzzo Joseph, R. Levine Diana, K. Saeid Kamal, C. Schwartz Rachel, T. Sucic Nicholas, D. Terleckyj Oksana, M. Young Jeffrey, Bacterial communication ("quorum sensing") via ligands and receptors: a novel pharmacologic target for the design of antibiotic drugs. *J. Pharmacol. Exp. Ther*, 312(2) 2005, 417-23.
- [9] T.B. Rasmussen, M. Givskov, Quorum-sensing inhibitors as antipathogenic drugs. *Int. J. Med. Microbiol*, 296 (2-3) 2006, 149-61.
- [10] B. Rasmussen Thomas, M. Givskov, Quorum sensing inhibitors: a bargain of effects. *Microbiology*, 152 (Pt 4) 2006, 895-904.
- [11] N. Balaban, M. Givskov, T.B. Rasmussen, In Vivo Studies: Inhibiting Biofilm-Associated Bacterial Infections using QSIs. In *Control of Biofilm Infections by Signal Manipulation*, Balaban, N. Ed. Springer: Berlin, 2 2008, 119-29.
- [12] D. Ren, M. Givskov, T. Ramussen, Quorum-Sensing Inhibitory Compounds. In *Control of Biofilm Infections by Signal Manipulation*, Balaban, N. Ed. Springer: Berlin, 2 2008, 51-77.
- [13] M. Hentzer, M. Givskov, L. Eberl, Quorum sensing in biofilms: Gossip in slime city. *Microb. Biofilms*, 1 2004, 118-40.
- [14] C.D. Nadell, J.B. Xavier, S.A. Levin, K.R. Foster, The evolution of quorum sensing in bacterial biofilms. *PLoS Biol*, 6 (1) 2008, 171-9.
- [15] M.R. Parsek, E.P. Greenberg, Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends Microbiol*, 13 (1) 2005, 27-33.
- [16] A.R. Horswill, P. Stoodley, P.S. Stewart, M.R. Parsek, The effect of the chemical, biological, and physical environment on quorum sensing in structured microbial communities. *Anal. Bioanal. Chem*, 387 (2) 2007, 371-80.
- [17] M. Li, N. Ni, H.T. Chou, C.D. Lu, P.C. Tai, B. Wang, Structure-based discovery and experimental verification of novel AI-2 quorum sensing inhibitors against *Vibrio harveyi*. *Chem Med Chem*, 3 (8) 2008, 1242-49.
- [18] J.B. Andersen, A. Heydorn, M. Hentzer, L. Eberl, O. Geisenberger, B.B. Christensen, S. Molin, M. Givskov, gfp-based N-acyl homoserine-lactone sensor systems for detection of bacterial communication. *Appl Environ Microbiol* 67 2001, 575-585.
- [19] W. Galloway, J. Hodgkinson, S. Bowden, M. Welch, D. Spring, Quorum Sensing in Gram-Negative Bacteria: Small-Molecule Modulation of AHL and AI-2 Quorum Sensing Pathways. *Chem. Rev*. 111 2011, 28-67.

- [20] K.M. Gray, J. Garey, The evolution of bacterial LuxI and LuxR quorum sensing regulators. *Microbiology*, 147 2001, 2379–2387.
- [21] S. Jafra, J.M. van der Wolf, Fast screening method for detection of acyl-HSL-degrading soil isolates. *J Microbiol Methods*, 57 2004, 415-420.
- [22] R. McLean, L. Pierson, C.A. Fuqua, simple screening protocol for the identification of quorum signal antagonists. *J. Microbiol. Methods*, 58 2004, 351-60.
- [23] T. Rasmussen, T. Bjarnsholt, M. Skindersoe, M. Hentzer, P. Kristoffersen, M. Koete, J. Nielsen, L. Eberl, M. Givskov, Screening for quorum-sensing inhibitors (QSI) by use of a novel genetic system, the QSI selector. *J. Bacteriol*, 187(5) 2005, 1799-1814.
- [24] M.G. Kocielek, Quorum-Sensing Inhibitors and Biofilms. *Anti-Infective Agents in Medicinal Chemistry*, 8 2009, 315-326 315.
- [25] L. Steindler, V. Venturi, Detection of quorum-sensing N-acyl homo serine lactone signal molecules by bacterial biosensors. *FEMS Microbiol. Lett*, 266 2007, 1-9.
- [26] C. Mendez, J. Salas, The role of ABC transporters in antibiotic-producing organisms: drug secretion and resistance mechanisms. *Research in Microbiology*, 152 2001, 341-350.
- [27] W. Bauer, M. Tepletski, Can plants manipulate bacterial quorum sensing? *Australian Journal of Plant Physiology*, 28 2001, 913-921.
- [28] S. Schauder, B. Bassler, The languages of bacteria. *Genes and Development*, 15 2001, 1468-1480.
- [29] R. Smith, B. Iglewski, *Pseudomonas aeruginosa* quorum-sensing systems and virulence. *Current Opinion in Microbiology*, 6 2003, 56-60.
- [30] H. Wu, Z. Song, M. Hentzer, J. Andersen, S. Molin, M. Givskov, N. Hoiby, Synthetic furanones inhibit quorum-sensing and enhance bacterial clearance in *Pseudomonas aeruginosa* lung infection in mice. *Journal of Antimicrobial Chemotherapy*, 53 2004, 1054-1061.
- [31] D. Cvitkovitch, Y. Liu, R. Ellen, Quorum sensing and biofilm formation in Streptococcal infections. *J Clin Invest*, 12 2003, 1626-1632.
- [32] E.P. Greenberg, Bacterial communication and group behavior. *J Clin Invest*, 112 2003, 1288-1290.
- [33] J.M. Yarwood, P.M. Schlievert, Quorum sensing in Staphylococcus infections. *J Clin Invest*, 112 2003, 1620-1625.
- [34] C. Arevalo-Ferro, J. Buschmann, G. Reil, A. Gorg, L. Wiehlmann, B. Tummeler, L. Eberl, K. Riedel, Proteome analysis of intracolonial diversity of two *Pseudomonas aeruginosa* TB clone isolates. *Proteomics* 4 2004, 1241-1246.
- [35] J.S Webb, M. Givskov, S. Kjelleberg, Bacterial biofilms: prokaryotic adventures in multicellularity. *Curr Opin Microbiol*, 6 2003, 578-585.
- [36] J. Zhu, J. Mekalanos, Quorum sensing-dependent biofilms enhance colonization in *Vibrio cholerae*. *Dev Cell* 5 2003, 647-656.
- [37] J.M. Dow, L. Crossman, K. Findlay, Y. He, J. Feng, J. Tang, Biofilm dispersal in *Xanthomonas campestris* is controlled by cell-cell signaling and is required for full virulence to plants. *Proc Natl Acad Sci USA*, 100 2003, 10995-11000.
- [38] K.B. Xavier, B. Bassler, LuxS quorum sensing: more than just a numbers game. *Curr Opin Microbiol*, 6 2003, 191-197.
- [39] V. Sperandio, A. Torres, B. Jarvis, J. Nataro, J. Kaper, Bacteria-host communication: The language of hormones. *Proc Natl Acad Sci USA*, 100 2003, 8951-8956.
- [40] S.C. Williams, E. Patterson, N. Carty, J. Griswold, A. Hamood, K. Rumbaugh, *Pseudomonas aeruginosa* autoinducer enters and functions in mammalian cells. *J Bacteriol*, 186 2004, 2281-2287.
- [41] B. Stevenson, K. von Lackum, R. Wattier, J. McAlister, J. Miller, K. Babb, Quorum sensing by the Lyme disease spirochete. *Microbes Infect*, 5 2003, 991-997.
- [42] D. Ren, T.K. Wood, (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone reduces corrosion from *Desulfotomaculum orientis*. *Environ. Microbiol*, 6(5) 2004, 535-40.
- [43] T. Rasmussen, M. Manefield, J. Andersen, L. Eberl, U. Anthoni, C. Christophersen, P. Steinberg, S. Kjelleberg, M. Givskov, How *Delisea pulchra* furanones affect quorum sensing and swarming motility in *Serratia liquefaciens* MG1. *Microbiology*, 146(12) 2000, 146, 3237-44.
- [44] J.M. Henke, B. Bassler, Bacterial social engagements. *Trends in Cell Biology*, 14 2004, 648-656.

- [45] N.A. Whitehead, A. Barnard, H. Slater, Simpson, N.J., Salmond, G.P. Quorum-sensing in Gram-negative bacteria. *FEMS Microbiology Reviews*, 25 2001, 365-404.
- [46] N. Halliday, K. Hardie, P. Williams, K. Winzer, D. Barrett, Quantitative liquid chromatography-tandem mass spectrometry profiling of activated methyl cycle metabolites involved in LuxS-dependent quorum sensing in *Escherichia coli*. *Anal Biochem*, 403 2010, 1-2.
- [47] S. Hagen, M. Son, J. Weiss, J. Young, Bacterium in a box: sensing of quorum and environment by the LuxI/LuxR gene regulatory circuit. *J Biol Phys*, 36 2010, 317-27.
- [48] V Sperandio, A. Sdi, Sensing of acyl-homoserine lactones by enterohemorrhagic *E. coli* (EHEC) serotype O157:H7 in the bovine rumen. *Gut Microbes*, 1 2010, 432-5.
- [49] M. Cao, Y. Feng, C. Wang, F. Zheng, M. Li, H. Liao, Y. Mao, X. Pan, J. Wang, D. Hu, F. Hu, J. Tang, Functional definition of LuxS, an autoinducer-2 (AI-2) synthase and its role in full virulence of *Streptococcus suis* serotype 2. *J Microbiol*, 49 2011, 1000-11.
- [50] S. Von Bodman, W. Bauer, D. Coplin, Quorum sensing in plant-pathogenic bacteria. *Annu Rev Phytopathol*, 41 2003, 455-482.
- [51] Y. Dong, L. Wang, L. Xu, H. Zhang, X. Zhang, L. Zhang, Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature*, 411 2001, 813-817.
- [52] C. Fuqua, E. Greenberg, Listening in on bacteria: acyl-homoserine lactone signalling. *Nature Reviews Molecular and Cellular Biology*, 3 2002, 685-695.
- [53] C. Fuqua, M. Parsek, E. Greenberg, Regulation of gene expression by cell-to-cell communication: Acyl-homoserine lactone quorum sensing. *Annu Rev Genet*, 35 2001, 439-468.
- [54] N. Decanis, N. Tazi, A. Correia, M. Vilanova, M. Rouabhia, A. Farnesol, a fungal quorum-sensing molecule triggers *Candida albicans* morphological changes by down regulating the expression of different secreted aspartyl proteinase genes. *Open Microbiol J*, 5 2011, 119-26.
- [55] K. Ohtani, H. Hayashi, T. Shimizu, The LuxS gene is involved in cell-cell signaling for toxin production in *Clostridium perfringens*. *Mol. Microbiol*, 44 2000, 171-179.
- [56] J. Smith, B. Ahmer, Detection of other microbial species by *Salmonella*: expression of the SdiA regulon. *J. Bacteriol*, 185 2003, 1357-1366.
- [57] E.W. Debler, G. Kaufmann, R. Kirchdoerfer, J. Mee, K. Janda, I. Wilson, Crystal structures of a quorum-quenching antibody. *Journal of Molecular Biology*, 368 2007, 1392-1402.
- [58] R. Smith, B. Iglewski, *Pseudomonas aeruginosa* quorum sensing as a potential antimicrobial target. *Journal of Clinical Investigation*, 112 2003, 1460-1465.
- [59] T. Hoang, S. Sullivan, J. Cusick, H. Schweizer, {beta}-Ketoacyl acyl carrier protein reductase (FabG) activity of the fatty acid biosynthetic pathway is a determining factor of 3-oxo-homoserine lactone acyl chain lengths. *Microbiology*, 148 2002, 3849-3856.
- [60] Y. Dong, L. Zhang, Quorum sensing and quorum-quenching enzymes. *Journal of Microbiology*, 43 2005, 101-109.
- [61] M. Schuster, E. Greenberg, A network of networks: quorum-sensing gene regulation in *Pseudomonas aeruginosa*. *Inter J Med Microbiol*, 296 2006, 73-81.
- [62] M. Hentzer, L. Eberl, J. Nielsen, M. Givskov, Quorum sensing-A novel target for the treatment of biofilm infections. *Bio Drugs*, 17 2003, 241-250.
- [63] J. Downie, The roles of extracellular proteins, polysaccharides and signals in the interactions of rhizobia with legume roots. *FEMS Microbiol Rev*, 34 2010, 150-70.
- [64] I. Khafagi, A. Dewedar, The efficiency of random versus ethno-directed research in the evaluation of Sinai medicinal plants for bioactive compounds. *Journal of Ethnopharmacology* 71 2000, 365-376.
- [65] N. Ni, H. Chou, J. Wang, M. Li, C. Lu, P. Tai, B. Wang, Identification of boronic acids as antagonists of bacterial quorum sensing in *Vibrio harveyi*. *Biochem. Biophys. Res. Commun*, 369(2) 2008, 590-94.
- [66] P. Williams, Quorum sensing, communication and cross-kingdom signalling in the bacterial world. *Microbiology*, 153(12) 2007, 3923-38.
- [67] M. Zahin, S. Hasan, F. Aqil, M. Khan, F. Husain, I. Ahmad, Screening of certain medicinal plants from India for their anti-quorum sensing activity. *Indian J Exp Biol*. 48 2010, 1219-24.

- [68] S. Schulz, J. Dickschat, B. Kunze, I. Wagner-Dobler, R. Diestel, F. Sasse, Biological activity of volatiles from marine and terrestrial bacteria. *Mar Drugs*, 8 2010, 2976-87.
- [69] H. Zhu, W. Liu, B. Tian, H. Liu, S. Ning, Inhibition of Quorum Sensing in the Opportunistic Pathogenic Bacterium *Chromobacterium violaceum* by an Extract from Fruiting Bodies of Lingzhi or Reishi Medicinal Mushroom, *Ganoderma lucidum* (W.Curt.:Fr.) P. Karst. (Higher Basidiomycetes). *Int J Med Mushrooms*. 13 2011, 559-64.
- [70] J.C. Espin, M. Garcia-Conesa, F. Tomas-Barberan, Nutraceuticals: facts and fiction. *Phytochemistry*, 68 2007, 2986-3008.
- [71] D. Vatter, K. Mihalik, S. Crixell, R. McLean, Dietary phytochemicals as quorum sensing inhibitors. *Fitoterapia*, 78 2007, 302-310.
- [72] B. Huber, L. Eberl, W. Feucht, J. Polster, Influence of polyphenols on bacterial biofilm formation and quorum-sensing. *Zeitschrift fur Naturforschung Section C, Biosciences*. 58 2004, 879-884.
- [73] G. Hernandez, J. Olmos, Molecular identification of pathogenic and nonpathogenic strains of *Vibrio harveyi* using PCR and RAPD. *Appl Microbiol Biotechnol*, 63 2004, 722-727.
- [74] K. Kim, Y. Kim, B. Koh, S. Hwang, S. Kim, F. Lepine, Y. Cho, G. Lee, HHQ and PQS, two *Pseudomonas aeruginosa* quorum-sensing molecules, down-regulate the innate immune responses through the nuclear factor-kappaB pathway. *Immunology*, 129 2010, 578-88.
- [75] Y. Wang C. Deng, Q. Peng, Z. Chen, D. Huang, J. Zhang, F. Song, Effect of quorum sensing response regulator nprR deletion on expression of cry protein in *Bacillus thuringiensis*. *Wei Sheng Wu Xue Bao*, 50 2010, 1550-1555.
- [76] L. Dițu, C. Chifiriuc, V. Lazar, G. Mihaescu, Implication of quorum sensing phenomenon in the expression of genes that code for bacteriocines in lactic bacteria. *Bacteriol Virusol Parazitol Epidemiol*, 54 2009, 147-66.
- [77] V. Lazar, Y. Miyazaki, T. Hanawa, M. Chifiriuc, M. Dițu, L. Maruțescu, C. Bleotu, S. Kamiya, The influence of some probiotic supernatants on the growth and virulence features expression of several selected enteroaggregative *E. coli* clinical strains. *Roum Arch Microbiol Immunol*, 68 2009 207-14.
- [78] C. Tsao, L. Wang, Y. Hashimoto, H. Yi, J. March, M. DeLisa, T. Wood, J. Valdes, W. Bentley, LuxS coexpression enhances yields of recombinant proteins in *Escherichia coli* in part through posttranscriptional control of GroEL. *Appl Environ Microbiol*, 77 2011, 2141-52.
- [79] M. Hentzer, M. Givskov, Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections. *J Clin Invest*, 112 2003, 1300-1307.
- [80] M. Manefield, L. Harris, S. Rice, R. De Nys, S. Kjelleberg, Inhibition of luminescence and virulence in the black tiger prawn (*Penaeus monodon*) pathogen *Vibrio harveyi* by intercellular signal antagonists. *Appl. Environ. Microbiol*, 66(5) 2000, 2079-84.
- [81] S. West, A. Griffin, A. Gardner, S. Diggle, Social evolution theory for microorganisms. *Nat Rev Microbiol*, 4 2006, 597-607.
- [82] K. Mok, N. Wingreen, B. Bassler, *Vibrio harveyi* quorum sensing: a coincidence detector for two autoinducers controls gene expression. *EMBO J*. 22 2003, 870-881.
- [83] M. Teplitski, J. Robinson, W. Bauer, Plants secrete substances that mimic bacterial N-acyl homo serine lactone signal activities and affect population density-dependent behaviors in associated bacteria. *Molecular Plant-Microbe Interactions*. 13 2000, 637-648.
- [84] M. Gao, M. Teplitski, J. Robinson, W. Bauer, Production of substances by *Medicago truncatula* that affect bacterial quorum sensing. *Mol. Plant Microbe Interact*, 16 2003, 827-834.
- [85] A. Adonizio, K. Kong, K. Mathee, Inhibition of quorum sensing-controlled virulence factor production in *Pseudomonas aeruginosa* by south Florida plant extracts. *Antimicrob. Agents Chemother*, 52(1) 2008, 198-203.
- [86] D. Hogan, A. Vik, R. Kolter. A *Pseudomonas aeruginosa* quorum-sensing molecule influences *Candida albicans* morphology. *Mol. Microbiol*, 54 2004, 1212-1223.
- [87] T. Persson, H. Hansen Thomas, B. Rasmussen Thomas, E. Skinderso Mette, M. Givskov, J. Nielsen, J. Rational design and synthesis of new quorum-sensing inhibitors derived from acylated

- homoserine lactones and natural products from garlic. *Org. Biomol. Chem*, 3(2) 2005, 253-62.
- [88] T. Rasmussen, M. Skindersoe, T. Bjarnsholt, R. Phipps, K. Christensen, P. Jensen, J. Andersen, J. B. Koch, T. Larsen, M. Hentzer, L. Eberl, N. Hoiby, M. Givskov, Identity and effects of quorum-sensing inhibitors produced by *Penicillium* species. *Microbiology*, 151(5) 2005, 1325-40.
- [89] G. Nascimento, P. Locatelli, P. Freitas, G. Silva, Antimicrobial activity of plant extracts and phytochemicals on antibiotic resistant bacteria. *Braz. J. Microbiol*, 31 2000, 247-256
- [90] R. Al-Hussaini, A. Mahasneh, Microbial Growth and Quorum Sensing Antagonist Activities of Herbal Plants Extracts. *Molecules*. 14 2009, 3425-3435.
- [91] C. Boga, L. Forlani, R. Calienni, T. Hindley, A. Hochkoepler, S. Tozzi, N. Zanna, on the antibacterial activity of roots of *Capparis spinosa* L. *Nat Prod Res*, 25 2011, 417-421.
- [92] T. Krishnan, W. Wai-Fong, Yin, K. Chan, Inhibition of Quorum Sensing Controlled Virulence Factor Production in *Pseudomonas aeruginosa* PAO1 by Ayurveda Spice Clove (*Syzygium Aromaticum*) Bud Extract. *Sensors*, 12 2012, 4016-4030.
- [93] S. Issac Abraham, A. Palani, B. Ramaswamy, K. Shunmugiah, V. Arumugam, Antiquorum sensing and antibiofilm potential of *Capparis spinosa*. *Arch Med Res*, 42 2011, 658-668.
- [94] Y. Chong, W. Yin, C. Ho, M. Mustafa, A. Hadi, K. Awang, P. Narrima, C. Koh, Malabaricone C from *Myristica Cinnamomea* exhibits Anti-Quorum sensing activity. *Journal of Natural Products*, 74 2011, 2261-2264.
- [95] B.N. Singh, H. Singh, A. Singh, B. Singh, A. Mishra, C. Nautiyal, *Lagerstroemia speciosa* fruit extract modulates quorum sensing-controlled virulence factor production and biofilm formation in *Pseudomonas. Aeruginosa*. *Microbiology*, 158 2012, 529-538.
- [96] L. Tan, W. Yin, K. Chan, Silencing Quorum Sensing through Extracts of *Melicope lunu-ankenda*. *Sensors*. 12 2012, 4339-4351.
- [97] S. Yeo, F. Tham, Anti-quorum sensing and antimicrobial activities of some traditional Chinese medicinal plants commonly used in South-East Asia. *Malaysian Journal of Microbiol*, 8 2012, 11-20.
- [98] S. Chun, D. Vattem, Y. Lin, K. Shetty, Phenolic antioxidants from clonal oregano (*Origanum vulgare*) with antimicrobial activity against *Helicobacter pylori*. *Process Biochemistry*, 40 2005, 809-816.
- [99] M. Rodrigues, L. Krause, E. Caramo, J. Dos Santos, C. Dariva, Vladimir de Oliveira, J. Chemical composition and extraction yield of the extract of *Origanum vulgare* obtained from sub- and supercritical CO₂. *Journal of Agricultural and Food Chemistry*, 52 2004, 3042-3047.
- [100] F. Chami, N. Chami, S. Bennis, J. Trouillas, A. Remmal, Evaluation of carvacrol and eugenol as prophylaxis and treatment of vaginal candidiasis in an immunosuppressed rat model. *Journal of Antimicrobial Chemotherapy*. 54 2004, 909-914.
- [101] P. Hersch-Martinez, B. Leanos-Miranda, F. Solórzano-Santos, Antibacterial effects of commercial essential oils over locally prevalent pathogenic strains in Mexico. *Fitoterapia*, 76 2005, 453-457.
- [102] M. Asif Saeed, A. Sabir, Antibacterial activities of some constituents from oleo-gum-resin of *Commiphora mukul*. *Fitoterapia*. 75 2004, 204-208.
- [103] B. Girenavar, M. Cepeda, K. Soni, A. Vikram, P. Jesudhasan, G. Jayaprakasha, S. Pillai, B. Patil, Grapefruit juice and its furocoumarins inhibits autoinducer signaling and biofilm formation in bacteria. *Int. J. Food Microbiol*, 125(2) 2008, 204-8.
- [104] B. Girenavar, G. Jayaprakasha, B. Patil, Potent inhibition of human cytochrome P450 3A4, 2D₆, and 2C₉ isoenzymes by grapefruit juice and its furocoumarins. *J. Food Sci*, 72(8) 2007, C417-21.
- [105] K. Lemar, M. Turner, D. Lloyd, Garlic (*Allium sativum*) as an anti-Candida agent: a comparison of the efficacy of fresh garlic and freeze-dried extracts. *J. Appl. Microbiol*, 93 2002, 398.
- [106] J. Shuford, J. Steckelberg, R. Patel, Effects of fresh garlic extract on candida albicans biofilms. *Antimicrob. Agents Chemother*, 49(1) 2005, 473.
- [107] T. Bjarnsholt, P. Jensen, T. Rasmussen, L. Christophersen, H. Calum, M. Hentzer, Garlic blocks quorum sensing and promotes rapid

- clearing of pulmonary *Pseudomonas aeruginosa* infections. *Microbiology*, 151 2005, 3873–3880.
- [108] J. Park, G. Kaufmann, J. Bowen, J. Arbiser, J. Janda, A. Solenopsis, a venom alkaloid from the fire ant *Solenopsis invicta*, inhibits quorum-sensing signaling in *Pseudomonas aeruginosa*. *J. Infect. Dis.*, 198(8) 2008, 1198-201.
- [109] J. Arbiser, T. Kau, M. Konar, K. Narra, R. Ramchandran, S. Summers, C. Vlahos, K. Ye, the alkaloidal component of the fire ant (*Solenopsis invicta*), is a naturally occurring inhibitor of phosphatidylinositol-3-kinase signaling and angiogenesis. *Blood*, 109(2) 2007, 560-65.
- [110] G. Howell, J. Butler, R. de Shazo, J. Farley, et al. Cardiodepressant and neurologic actions of *Solenopsis invicta* (imported fire ant) venom alkaloids. *Ann. Allergy, Asthma, Immunol*, 94(3) 2005, 380-86.
- [111] C. Niu, S. Afre, E. Gilbert, Subinhibitory concentrations of cinnamaldehyde interfere with quorum sensing. *Lett. Appl. Microbiol*, 43(5) 2006, 489-94.
- [112] C. Niu, E. Gilbert, Colorimetric method for identifying plant essential oil components that affect biofilm formation and structure. *Appl. Environ. Microbiol*, 70(12) 2004, 6951-56.
- [113] G. Brackman, T. Defoirdt, C. Miyamoto, P. Bossier, S. Van Calenbergh, H. Nelis, T. Coenye, Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. By decreasing the DNA-binding activity of the quorum sensing response regulator LuxR. *BMC Microbiol*, 8 2008, 149.
- [114] J. Choo, Y. Rukayadi, J. Hwang, Inhibition of bacterial quorum sensing by vanilla extract. *Lett. Appl. Microbiol*, 42(6) 2006, 637-41.
- [115] A. Adonizio, K. Downum, B. Bennett. K. Mathee, Antiquorum sensing activity of medicinal plants in southern Florida. *J. Ethnopharmacol*, 103 2006, 427–435.
- [116] B. Huber, L. Eberl, W. Feucht, J. Polster, Influence of Polyphenols on Bacterial Biofilm Formation and Quorum-sensing. *Z. Naturforsch.* 58 2003, 879-884.
- [117] N. Ni, G. Choudhary, M. Li, B. Wang, Pyrogallol and its analogs can antagonize bacterial quorum sensing in *Vibrio harveyi*. *Bioorg. Med. Chem. Lett.*, 18(5) 2008, 1567-72.
- [118] Z. Zeng, L. Qian, L. Cao, Virtual screening for novel quorum sensing inhibitors to eradicate biofilm formation of *Pseudomonas aeruginosa*. *Appl. Microbiol. Biotechnol*, 79(1) 2008, 119-26.
- [119] T. Walker, H. Bais, E. Deziel, H. Schweizer, *Pseudomonas aeruginosa*-plant root interactions. Pathogenicity, biofilm formation, and root exudation. *Plant Physiol*, 134 2004, 320-331.
- [120] M. Otto, Quorum-sensing control in *Staphylococci* - a target for antimicrobial drug therapy? *FEMS Microbiol. Lett*, 241 2004, 24(2) 135-41.
- [121] N. Balaban, L. Collins, J. Cullor, E. Hume, Medina-Acosta, E, Da Motta, O. V, O'Callaghan, R, Rossitto, P. V, Shirliff, M. E, Da Silveira, L. S, Tarkowski, A, Torres, J. V. *Prevention of diseases caused by Staphylococcus aureus using the peptide RIP. Peptides*, 21(9) 2000, 1301-11.
- [122] Y. Gov, A. Bitler, G. Dell'Acqua, RNAIII inhibiting peptide (RIP), a global inhibitor of *Staphylococcus aureus* pathogenesis: structure and function analysis. *Peptides (N. Y. NY, USA)*, 22(10) 2001, 1609-20.
- [123] N. Balaban, O. Cirioni, A. Giacometti, Treatment of *Staphylococcus aureus* biofilm infection by the quorum-sensing inhibitor RIP. *Antimicrob. Agents Chemother*, 51(6) 2007, 2226-9.
- [124] N. Balaban, A. Giacometti, O. Cirioni, Use of the quorum-sensing inhibitor RNAIIIinhibiting peptide to prevent biofilm formation in vivo by drugresistant *Staphylococcus epidermidis*. *J. Infect. Dis.*, 187(4) 2003, 625-30.
- [125] O. Cirioni, R. Ghiselli, D. Minardi, RNAIII-inhibiting peptide affects biofilm formation in a rat model of *Staphylococcal* ureteral stent infection. *Antimicrob. Agents Chemother*, 51(12) 2007, 4518-20.
- [126] G. Dell'Acqua, A. Giacometti, O. Cirioni, Suppression of drug-resistant staphylococcal infections by the quorum-sensing inhibitor RNA III inhibiting peptide. *J. Infect. Dis.*, 190(2) 2004, 318-20.
- [127] P. Anguita-Alonso, A. Giacometti, RNAIII-inhibiting-peptide-loaded olymethacrylate prevents in vivo *Staphylococcus aureus* biofilm formation. *Antimicrob. Agents Chemother*, 51(7), 2007, 2594-96.

- [128] N. Balaban, Y. Gov, A chimeric peptide composed of a dermaseptin derivative and an RNA III-inhibiting peptide prevents graft-associated infections by antibiotic-resistant staphylococci. *Antimicrob. Agents Chemother*, 48(7) 2004, 2544-50.
- [129] O. Cirioni, A. Giacometti, R. Ghiselli, RNAIII-inhibiting peptide significantly reduces bacterial load and enhances the effect of antibiotics in the treatment of central venous catheter-associated Staphylococcus aureus infections. *J. Infect. Dis*, 193(2) 2006, 180-86.
- [130] P. Domenico, E. Gurzenda, A. Giacometti, Bis EDT and RIP act in synergy to prevent graft infections by resistant staphylococci. *Peptides*, 25(12) 2004, 2047-53.
- [131] A. Giacometti, O. Cirioni, R. Ghiselli, G. Dell'Acqua, RNAIII-inhibiting peptide improves efficacy of clinically used antibiotics in a murine model of staphylococcal sepsis. *Peptides*, 2005, 26(2), 169-75.
- [132] Chan, W. C, Coyle, B. J, Williams, P. Virulence Regulation and Quorum Sensing in Staphylococcal Infections: Competitive AgrC Antagonists as Quorum Sensing Inhibitors. *J. Med. Chem*, 47(19) 2004, 4633-41.
- [133] G. Lyon, P. Mayville, T. Muir, Rational design of a global inhibitor of the virulence response in Staphylococcus aureus, based in part on localization of the site of inhibition to the receptor-histidine kinase, *AgrC*. *Proc. Natl. Acad. Sci. USA*, 97(24) 2000, 13330-35.
- [134] G. Lyon, T. Muir, Chemical signaling among bacteria and its inhibition. *Chem Biol*, 10 2003, 1007-1021.
- [135] J. Wright, R. III, Jin, R. Novick, R. Transient interference with staphylococcal quorum sensing blocks abscess formation. *Proc. Natl. Acad. Sci. USA*, 102(5) 2005, 1691-96.
- [136] J. Wright, G. III, Lyon, E. George, Hydrophobic interactions drive ligand-receptor recognition for activation and inhibition of staphylococcal quorum sensing. *Proc. Natl. Acad. Sci. USA*, 101(46) 2004, 16168-73.
- [137] K. Tateda, R. Comte, J. Pechere, T. Kohler, Azithromycin inhibits quorum sensing in Pseudomonas aeruginosa. *Antimicrob. Agents Chemother*, 45(6) 2001, 1930-33.
- [138] M. Skindersoe, M. Alhede, R. Phipps, Effects of antibiotics on quorum sensing in Pseudomonas aeruginosa. *Antimicrob. Agents Chemother*, 52(10) 2008, 3648-63.
- [139] S. Favre-Bonte, T. Koehler, C. Van Delden, Biofilm formation by Pseudomonas aeruginosa: role of the C4-HSL cell-to-cell signal and inhibition by azithromycin. *J. Antimicrob. Chemother*, 52(4) 2003, 598-604.
- [140] N. Hoffmann, B. Lee, Azithromycin blocks quorum sensing and alginate polymer formation and increases the sensitivity to serum and stationary-growth-phase killing of Pseudomonas aeruginosa and attenuates chronic P. aeruginosa lung infection in Cfr-/- mice. *Antimicrob. Agents Chemother*, 51(10) 2007, 3677-87.
- [141] Y. Yoon, J. Sofos, Absence of association of autoinducer-2-based quorum sensing with heat and acid resistance of Salmonella. *J Food Sci*, 75 2010, 444-448.
- [142] D. Martinelli, G. Grossmann, Effects of natural and chemically synthesized furanones on quorum sensing in Chromobacterium violaceum. *BMC Microbiol*, 4 2004, 25-36.
- [143] D. Ren, J. Sims, T. Wood, Inhibition of biofilm formation and swarming of Escherichia coli by (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone. *Environ. Microbiol*, 3(11) 2001, 731-6.
- [144] D. Ren, L. Bedzyk, P. Setlow, Differential gene expression to investigate the effect of (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone on Bacillus subtilis. *Appl. Environ. Microbiol*, 70(8) 2004, 4941-49.
- [145] D. Ren, J. Sims, T. Wood, Inhibition of biofilm formation and swarming of Bacillus subtilis by (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone. *Lett. Appl. Microbiol*, 34(4) 2002, 293-99.
- [146] D. Ren, L. Bedzyk, R. Ye, Stationary-phase quorum-sensing signals affect autoinducer-2 and gene expression in Escherichia coli. *Appl. Environ. Microbiol*, 70(4) 2004, 2038-43.
- [147] T. Benneche, Z. Hussain, A. Aamdal Scheie, Synthesis of 5-(bromomethylene)furan-2(5H)-ones and 3-(bromomethylene)isobenzofuran-1(3H)-ones as inhibitors of microbial quorum sensing. *New J. Chem*, 32(9) 2008, 1567-72.

- [148] K. Mathee, A. Adonizio, F. Ausubel, Use of Ellagitannins as Inhibitors of Bacterial Quorum Sensing. *PCT Int. Appl.* WO/2009/114810, 2009.
- [149] H. Liu, K. Koh, J. Kim, Y. Seo, *Biotechnol. Bioprocess Eng.* 13 2008, 458-466.
- [150] K. Stephenson, J. Hoch, Developing inhibitors to selectively target two-component and phosphorelay signal transduction systems of pathogenic microorganisms. *Curr Med Chem*, 11 2004, 765-773.
- [151] G. Lyon, J. Wright, T. Muir, R. Novick, Key Determinants of Receptor Activation in the agr Autoinducing Peptides of *Staphylococcus aureus*. *Biochemistry*, 41(31) 2002, 10095-104.
- [152] V. Denervaud, P. TuQuoc, D. Blanc, S. Favre-Bonte, V. Krishnapillai, Characterization of cell-to-cell signaling-deficient *Pseudomonas aeruginosa* strains colonizing intubated patients. *J Clin Microbiol*, 42 2004, 554-562.
- [153] K. Duan, C. Dammel J. Stein H. Rabin, Surette MG: Modulation of *Pseudomonas aeruginosa* gene expression by hostmicroflora through interspecies communication. *Mol Microbiol*, 50 2003, 1477-1491.
- [154] A. Bozorgian, S. Zarinabadi, A. Samimi, Preparation of Xanthan Magnetic Biocompatible Nano-Composite for Removal of Ni²⁺ from Aqueous Solution, *Chemical Methodologies*, 4 (4) (2020), 477-493
- [155] A. Samimi, S. Zarinabadi, A. Bozorgian, A. Amosoltani, M. Tarkesh, K. Kavousi, Advances of Membrane Technology in Acid Gas Removal in Industries, *Progress in Chemical and Biochemical Research*, 3 (1) (2020), 46-54
- [156] C. Aguilar, I. Bertani, V. Venturi, Quorum-sensing system and stationary-phase sigma factor (rpoS) of the onion pathogen *Burkholderia cepacia* genomovar I type strain, ATCC 25416. *Appl Environ Microbiol*, 69 2003, 1739-1747.
- [157] J. Hornby, S. Jacobitz-Kizzier, D. McNeel, Jensen, Inoculum size effect in dimorphic fungi: extracellular control of yeast-mycelium dimorphism in *Ceratocystis ulmi*. *Appl Environ Microbiol*, 70 2004, 1356-1359.
- [158] U. Mathesius, S. Mulders, S. Gao, M. Teplitski, G. Caetano-Anolles, B. Rolfe, Extensive and specific responses of a eukaryote to bacterial quorum-sensing signals. *Proc Natl Acad Sci USA*, 100 2003, 1444-1449.
- [159] G. Fleet, Yeast interactions and wine flavour. *Int J Food Microbiol*, 86 2003, 11-22.
- [160] Y. Dong, A. Gusti, Q. Zhang, Identification of quorum-quenching N-acyl homoserine lactonases from *Bacillus* species. *Appl Environ Microbiol*, 68 2002, 1754-1759.
- [161] L. Molina, F. Constantinescu, Degradation of pathogen quorum-sensing molecules by soil bacteria: a preventive and curative biological control mechanism. *FEMS Microbiol Ecol*, 45 2003, 71-81.
- [162] F. Xu, T. Byun, Degradation of Nacylhomoserine lactones, the bacterial quorum-sensing molecules, by acylase. *J Biotechnol*, 101 2003, 89-96.
- [163] M. Pasmore, J.W. Costerton, Biofilms, bacterial signaling, and their ties to marine biology. *J Ind Microbiol Biotechnol*, 30 2003, 407-413.
- [164] J. Skerratt, J. Bowman, G. Hallegraeff, S. James, Algicidal bacteria associated with blooms of a toxic dinoflagellate in a temperate Australian estuary. *Mar Ecol Prog Ser*, 244 2002, 1-15.
- [165] A. Valle, M. Bailey, A. Whiteley, N-acyl-Lhomoserine lactones (AHLs) affect microbial community composition and function in activated sludge. *Environ Microbiol*, 6 2004, 424-433.
- [166] R. Van Houdt, A. Aertsen, A. Jansen, A. Quintana, Biofilm formation and cell-to-cell signalling in Gram-negative bacteria isolated from a food processing environment. *J Appl Microbiol*, 96 2004, 177-184.
- [167] C. Waters, B. Bassler, Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol*, 21 2005, 319-346.
- [168] J. Njoroge, V. Sperandio, Jamming bacterial communication: new approaches for the treatment of infectious diseases. *EMBO Mol Med*, 1 2009, 201-10.
- [169] P. Ronald, Small protein-mediated quorum sensing in a Gram-negative bacterium: novel targets for control of infectious disease. *Discov Med*, 12 2011, 461-70.
- [170] T. Czarán, Janus-headed communication promotes bacterial cooperation and cheating: is quorum sensing useful against infections? *Virulence*, 1 2010, 402-403.
- [171] Y. Dong X. Zhang, Insecticidal *Bacillus thuringiensis* silences *Erwinia carotovora*

- virulence by a new form of microbial antagonism, signal interference. *Appl Environ Microbiol*, 70 2004, 954-960.
- [172] R. Dudler, L. Eberl, Interactions between bacteria and eukaryotes via small molecules. *Curr. Opin. Biotechnol*, 17 2006, 268–273.
- [173] J. March, W. Bentley, Quorum sensing and bacterial cross-talk in biotechnology. *Current Opinion in Biotechnology*, 15 2004, 495–502.
- [174] Z. Hosseinzadeh, A. Ramazani, N. Razzaghi-Asl, Plants of the Genus *Heracleum* as a Source of Coumarin and Furanocoumarin, *Journal of chemical reviews*, 1(2) 2020, 78-98.
- [175] M. Abdollahbeigi, Optimizing the Process of Di-Isobutyl Phthalate Production Using Vapor Permeation, *DAV International Journal of Science*, 4(2) (2015), 47-52
- [176] M.J. Choobineh, B. Nasrollahzadeh, M. Abdollahbeigi, Investigation of Contact Resistance Effect on Finned Pipes under Natural and Forced Convection, *DAV International Journal of Science*, 4(2) (2015), 58-76
- [177] B. Nasrollahzadeh, M.J. Choobineh, M. Abdollahbeigi, Investigation of Hydrate Formation Kinetics and Mechanism of Inhibitors Effect, *DAV International Journal of Science*, 4 (2015), 49-56
- [178] M. Abdollahbeigi, M.J. Choobineh, B. Nasrollahzadeh, Nano Catalyst, Operation Mechanism and Their Application in Industry, *Australian Journal of International Social Research*, 1(5) (2015), 1-6
- [179] M. Abdollahbeigi, M.J. Choobineh, B. Nasrollahzadeh, Investigation of Molecular Structure in Gas Hydrate, *Science road Journal*, 3(12) (2015), 74-79
- [180] M.J. Choobineh, M. Abdollahbeigi, B. Nasrollahzadeh, the formation of gas hydrate and the effect of inhibitors on their formation process, *Journal of Fundamental Applied Science* 8(2S) (2016), 1150-1159
- [181] A. Bozorgian, S. Zarinabadi, A. Samimi, Optimization of Well Production by Designing a Core pipe in one of the Southwest oil Wells of Iran, *Journal of Chemical Reviews*, 2(2) (2020), 122-129
- [182] D.Ren, A. Bedzyk Laura, W. Ye Rick, Differential gene expression shows natural brominated furanones interfere with the autoinducer-2 bacterial signaling system of *Escherichia coli*. *Biotechnol. Bioeng*, 88(5) 2004, 630-42.

How to cite this manuscript: Mohammad Asif, Quorum Sensing Inhibitors: Current Progress of Natural Antimicrobials, *Journal of Chemical Reviews (J. Chem. Rev.)*, 2020, 3(1), 320-339. DOI: 10.33945/SAMI/jcr.2019.174777.1020